

HOME GROUNDS FACT SHEET



Cornell University
Cooperative Extension
Nassau County



Horticulture Center
Demonstration & Community Gardens
at East Meadow Farm
832 Merrick Avenue
East Meadow, NY 11554
Phone: 516-565-5265

Maple Decline

by Juliet E. Carroll, Extension Plant Pathologist

Maple decline affects primarily sugar maple (*Acer saccharum*), Norway maple (*Acer platanoides*) and red maple (*Acer rubrum*) in the Northeast. The problem is not a new one; stagheaded maples were described as early as 1917 in Massachusetts. At that time, dieback was attributed mainly to drought and to the poor conditions for tree growth afforded by the urban environment. However, reports of the incidence and severity of maple decline have increased markedly in recent years to include urban, sugarbush and forest environments.

In forests, maples usually begin to decline after several successive years of defoliation by insects. Affected trees not only lose their first set of leaves to these insects, but will often use up valuable food reserves to produce a second set. During and after "refoliation," chemical changes occur in the tree that increases its susceptibility to secondary pathogens. *Armillaria mellea* (root rot), *Nectria cinnabarina* (branch canker) and *Steganosporium ovatum* (twig blight) are three fungi that frequently attack and may kill trees weakened by defoliation and refoliation.

In sugarbushes, predisposing stresses include drought, heavy grazing, over-tapping and/or heavy traffic by farm machinery. Seriously affected trees are often over-mature and have been heavily tapped for many years. Tapping holes, animal-damaged roots, and machine-damaged roots are all routes for entry of wood decay organisms. If this scenario is followed by insect defoliation as previously described, the result is often mortality of the stressed trees.

In urban sites principal stress factors in maple decline include drought, de-icing salts and/or road and sidewalk construction. These stresses also facilitate invasion by secondary organisms including root rots, decays and twig blights which greatly reduce chances of recovery from original stress(es). No matter which of the three environments maple decline occurs in, the sequence of events is similar. Healthy trees are stressed repeatedly, the stresses alter the trees' internal chemistry to allow repeated attack by secondary organisms, and the trees ultimately die.



Symptoms:

1. Reduced twig growth. Yearly twig growth varies considerably between trees and even within the canopies of individual trees. If the distance from bud scar to bud scar is less than or equal to five cm on a non-shaded twig, the tree may be in trouble.
2. Reduced foliage growth. Keep in mind the normal, healthy appearance of the particular maple species' foliage. Foliage that is sparse, light green and/or scorched signals that the tree may be declining.
3. Early fall coloration. Maples normally begin showing fall color after the first frost or in mid-to-late September. When fall color develops earlier than normal, in late July or early August, the maple is suffering from decline.


E-2-8 DWM/AAH revised RT 1/03

-continued-

Building Strong and Vibrant New York Communities

Cornell Cooperative Extension in Nassau County provides equal program and employment opportunities.

4. Dead branches in upper canopy. Small dead branches seen in tree tops in late spring or early summer are indicative of decline. Over time, larger, more visible branches and limbs will dieback. The more numerous the dead twigs or branches are, the more severe the decline condition.
5. Poor root condition. If roots can be examined, look for reduced occurrence of small feeder rootlets; dead, brittle roots; and decaying buttress roots.



Management:

Treatment for declining urban maples includes watering, fertilizing, pruning dead branches and reducing salt-laden spring water runoff over the roots. Thoroughly water trees every week or two during extended dry weather.

Use a soaker hose [a round, black, leaky (sweating) hose *or* flat green hose with a white stripe on the side]. If using the flat green hose, place holes facing down.

Place hose around the tree beyond the drip-line (outer foliage) and work in toward the trunk in a spiral fashion, to deliver water equally to all of the tree's roots.

Fertilize the trees with a complete fertilizer in the late fall or spring. The general recommendation is 2 to 4 lbs. of nitrogen per 1,000 sq. ft. of root area, or 1/4 to 1/2 lb. nitrogen per inch of tree diameter at breast height (measured with calipers). Broadcast the fertilizer over the surface of the ground. Some risk of burn on nearby turf may occur at the higher rates. For more information see Home Grounds Fact Sheet D-1-19, Fertilization of Trees and Shrubs.

Prune dead branches to possibly stimulate renewed, vigorous shoot growth. Pruning is best done in March, prior to budbreak, when the tree's ability to defend itself against disease is at its best. Road salt impact can be reduced by placing a barrier (curb, berm, ditch, etc.) which will catch and/or divert the spring runoff water which often contains copious amounts of salt. If soil and foliar analyses have been run and high sodium or chloride concentrations were found, then leaching the soil with fresh water and applying gypsum to improve soil structure or texture may be useful.

By the time symptoms are noticed, the tree may be beyond being restored to its original splendor. However, at this time another tree may be planted which will eventually replace the declining maple. In this way, the newly planted tree will have a few years to grow prior to the removal of the declining maple. Plant young maple trees away from roads to avoid de-icing salt problems.

The success of treatment to declining maples depends primarily on early detection of maple decline, the health of the tree prior to treatment, and/or its ability to respond to treatment. Positive diagnosis will often depend on "on the site" examination or the amount of information obtainable from the person submitting a sample. It is advisable to have valuable trees examined by a Certified Consulting Arborist. However, the prescribed treatments of fertilizing, watering and pruning will not damage healthy trees.

For more information:

Home Grounds Fact Sheets

- D-1-22 – Girdling Roots
- E-2-6 – Leaf Scorch of Shade Trees
- E-2-7 – Verticillium Wilt
- D-1-7 – Pruning
- D-1-37 – How to Water a Tree
- D-1-19 – Fertilization of Trees and Shrubs
- D-1-23 – Mulches